

NOXIOUS WEEDS OF GLACIER HILLS



Orange Hawkweed. Hairy leaves mostly found at base of plant. Stems mostly leafless with black bristly hairs.



stem and leaves.

Spotted Knapweed. Grayish-green



Spotted Knapweed



Spotted Knapweed (early)



Spotted Knapweed (early)



Canada Thistle



Canada Thistle. Prickly stems with stalkless leaves.



St. Johnswort. Yellow flowers with black dots at petal edge. Leaves with tiny transparent dots.



St. Johnswort (early)



Oxeye Daisy



Oxeye Daisy (early)



Oxeye Daisy (early)



Common Tansy (early). Dark green fern-like leaves, stems often purplish-red, yellow, button-like flowers in a dense, flat-topped cluster.



Common Tansy



Common Mullein



Common mullein. Fuzzy light green leaves thickly covered with soft, felt-like hairs. Single stout flowering stem up to 6 feet tall emerges in second year.



Leafy spurge. Heart-shaped floral leaves, greenish yellow flower clusters on long stalks cluster in flat-topped umbrellas.



Leafy spurge (early)



Leafy spurge



Meadow hawkweed



Meadow hawkweed (early)



Meadow hawkweed



Yellow toadflax



Yellow toadflax



Dalmatian toadflax



Houndstongue